

A Clinic Visit Case References

1. Liu C, Bayer A, Cosgrove SE, et al. Clinical practice guidelines by the infectious diseases society of america for the treatment of methicillin-resistant *Staphylococcus aureus* infections in adults and children. *Clin Infect Dis* 2011; 52:e18.
2. Stevens DL, Bisno AL, Chambers HF, et al. Practice guidelines for the diagnosis and management of skin and soft tissue infections: 2014 update by the infectious diseases society of America. *Clin Infect Dis* 2014; 59:147
3. Hepburn MJ, Dooley DP, Skidmore PJ, et al Comparison of short- course (5 days) and standard (10 days) treatment for uncomplicated cellulitis. *Arch Intern Med* 2004; 164:1669–74.
4. Rajendran PM, Young D, Maurer T, et al. Randomized, double-blind, placebo-controlled trial of cephalexin for treatment of uncomplicated skin abscesses in a population at risk for community-acquired methicillin-resistant *Staphylococcus aureus* infection. *Antimicrob Agents Chemother* 2007; 51:4044-8.
5. Schmitz GR, Bruner D, Pitotti R, et al. Randomized controlled trial of trimethoprim-sulfamethoxazole for uncomplicated skin abscesses in patients at risk for community-associated methicillin-resistant *Staphylococcus aureus* infection. *Ann Emerg Med* 2010; 56:283-7. 6. Talan DA et al. Trimethoprim-Sulfamethoxazole versus Placebo for Uncomplicated Skin Abscess. *N Engl J Med*. 2016; 374(9):823.